

# Calf BaseLyte™

## Electrolyte / Nutritional Supplement for Calves

### RECOMMENDED USES:

Nutritional supplement for dehydrated calves at shipping, during fluid loss, when experiencing intestinal upset, off-feed due to shipping, handling, disease, or hot weather, to reduce shrink, in conjunction with veterinary therapy for diarrhea, other times listed nutrients, electrolytes and beneficial microbial cultures are desired.

### GUARANTEED ANALYSIS: (minimum per gram)

<b>Salt (minimum)</b>	5.80 %
<b>Salt (maximum)</b>	6.90 %
<b>Sodium (minimum)</b>	6.90 %
<b>Sodium (maximum)</b>	8.20 %
<b>Potassium</b>	2.44 %
<b>Live Microbial Count</b>	62.5 million CFU (Lactobacillus acidophilus, Lactobacillus plantarum, Enterococcus faecium, Lactobacillus casei, Bifidobacterium bifidum, Bifidobacterium infantis, Bifidobacterium longum, Lactobacillus lactis, Pediococcus acidilactici, Bacillus amyloliquefaciens)

**INGREDIENTS:** Dextrose, sodium acetate, sodium propionate (preservative), sodium chloride, potassium chloride, glycine, sodium silico aluminate, inulin, hydrolyzed yeast, dried whole milk, thiamine mononitrate, riboflavin, niacin, calcium pantothenate, pyridoxine hydrochloride, folic acid, biotin, vitamin B12 supplement, ascorbic acid, natural and artificial flavor, dried Bacillus amyloliquefaciens fermentation product, dried Lactobacillus acidophilus fermentation product, dried Bifidobacterium bifidum fermentation product, dried Bifidobacterium infantis fermentation product, dried Bifidobacterium longum fermentation product, dried Lactobacillus plantarum fermentation product, dried Enterococcus faecium fermentation product, dried Lactobacillus casei fermentation product, dried Lactobacillus lactis fermentation product, dried Pediococcus acidilactici fermentation product.

### FEEDING DIRECTIONS: **Level Scoop = 40 grams electrolyte powder.**

#### **Dehydrated calves, calves with diarrhea**

Mix 80 grams (2 scoops) electrolyte powder with warm water to make 2 liters electrolyte solution. Feed 2 liters of resulting electrolyte solution twice daily or as a drench or with a stomach tube for calves up to 250 lbs. body weight. For calves fed milk or milk replacer, separate milk and electrolyte solution feeding by at least 30 minutes or alternately feed milk or electrolyte solution at intervals spaced evenly throughout day.

#### **Daily feeding during hot weather**

mix at rate of 40 grams (one scoop) per gallon of drinking water to 80 grams (2 scoops) per 5 gallons drinking water.

Space feedings as evenly throughout day as possible and separate electrolyte feedings from milk / milk replacer feedings by at least 30 minutes to avoid overloading abomasum. Do not mix this electrolyte powder with milk or milk replacer. Consult veterinarian for proper drenching / tubing procedures. Discard any mixture not used immediately.

**STORAGE:** Keep closed tightly between uses. Store in cool, dry place.

### PRODUCT CODE

<b>GN-00440</b>	80 gram single feeding packets (20 packs / box)
<b>GN-00441</b>	5 kilogram (11 lb) pouch (4 / case) - 62 feedings per pouch
<b>GN-00442</b>	20 kilogram (44 lb) bag (1 / case) - 250 feedings per bag
<b>GN-00442</b>	20 kilogram (44 lb) bag (50 bags / pallet)



TS\_CalfBaseLyteRevised10182021

Bio-Vet, Inc. • 300 Ernie Drive • Barneveld, WI 53507, U.S.A. • (800) 246- 8381 • Fax (608) 924- 7006 • www.Bio-Vet.com

Innovation at Work™